arXiv:1302.1274v1 [math.AC] 6 Feb 2013

TOP LOCAL COHOMOLOGY, MATLIS DUALITY AND TENSOR PRODUCTS

MAJID EGHBALI

ABSTRACT. Let $\mathfrak{a}$ be an ideal of a local ring $(R,\mathfrak{m})$ with $d = \dim R$. For $i > d$ the local cohomology module $H^i_{\mathfrak{a}}(R)$ vanishes. In the case that $H^d_{\mathfrak{a}}(R) \neq 0$, at first we give a bound for depth $D(H^d_{\mathfrak{a}}(R))$, where $d > 2$ and $(R,\mathfrak{m})$ is complete. More precisely we show

$$\operatorname{grade}(\mathfrak{b}, D(H^d_{\mathfrak{a}}(R))) + \dim R/\mathfrak{b} \geq \operatorname{depth}(D(H^d_{\mathfrak{a}}(R))) \geq 2,$$

where $\mathfrak{b}$ is an ideal of R of dimension ≤ 2 and D(-) denotes the Matlis duality functor. Later we give conditions that Cohen-Macaulayness of $D(H^d_{\mathfrak{a}}(R))$ fails, where $d \geq 3$. Finally, $H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R)$, $D(H^d_{\mathfrak{a}}(R)) \otimes_R D(H^d_{\mathfrak{a}}(R))$ and $H^d_{\mathfrak{a}}(R) \otimes_R D(H^d_{\mathfrak{a}}(R))$ are examined. In particular, we give the necessary and sufficient condition for Cohen-Macaulayness of $D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))$, where Q is a certain ideal of R.

1. INTRODUCTION

Let $\mathfrak{a} \subset R$ denote an ideal of a local ring $(R,\mathfrak{m})$ with $d = \dim R$. Let M be a finitely generated R-module. For an integer $i \in \mathbb{Z}$, let $H^i_{\mathfrak{a}}(M)$ denote the ith local cohomology module of M with respect to $\mathfrak{a}$ as introduced by Grothendieck (see [G] and [Br-Sh]). Let $E := E_R(R/\mathfrak{m})$ be the injective hull of the R-module $R/\mathfrak{m}$. By $D(-)$ we denote the Matlis dual functor $\operatorname{Hom}_R(-, E)$.

In algebraic geometry a (set-theoretic) complete intersection is a variety Y that can be cut out by $\operatorname{codim}(Y)$ many equations. By a result of Hellus [H, Corollary 1.1.4] Matlis dual of local cohomology can be useful to see whether we have a set-theoretic complete intersection ideal or not.

Of a particular interest is the last non-vanishing degree of the local cohomology module $H^i_{\mathfrak{a}}(R)$ known as the cohomological dimension $\operatorname{cd}(\mathfrak{a}, R)$. In general $\operatorname{cd}(\mathfrak{a}, R) \leq d$ and in the case that $\mathfrak{a} = \mathfrak{m}$, the equality occurs. Hartshorne-Lichtenbaum Vanishing Theorem gives the necessary and sufficient condition for non-vanishing of $H^d_{\mathfrak{a}}(R)$ (cf. [Br-Sh, Theorem 8.2.1]). Here we assume that $H^d_{\mathfrak{a}}(R) \neq 0$ and give some information on $D(H^d_{\mathfrak{a}}(R))$ and the tensor products of these modules. First, for a complete local ring $(R,\mathfrak{m})$ we show that depth $D(H^d_{\mathfrak{a}}(R)) \geq \min\{2, d\}$ (see Theorem 3.1). Then, we give an upper bound for $\operatorname{depth}(D(H^d_{\mathfrak{a}}(R)))$:

$$\operatorname{grade}(\mathfrak{b}, D(H^d_{\mathfrak{a}}(R))) + \dim R/\mathfrak{b} \geq \operatorname{depth}(D(H^d_{\mathfrak{a}}(R))) \geq 2,$$

where $\mathfrak{b}$ is an ideal of R of dimension at most 2 and $d \geq 3$. Hence, it provides conditions in which Cohen-Macaulayness of $D(H^d_{\mathfrak{a}}(R))$ fails ($d \geq 3$), see Remark 3.5.

Recently, there were some researches on the endomorphism ring of local cohomology modules and their importance, see for instance [H-St], [H-Sch], [Sch3] and [E-Sch]. However not so much is known on its dual, that is on the tensor product of local cohomology modules. Section four is devoted to examining the modules $H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R)$, $D(H^d_{\mathfrak{a}}(R)) \otimes_R D(H^d_{\mathfrak{a}}(R))$ and $H^d_{\mathfrak{a}}(R) \otimes_R$

2000 *Mathematics Subject Classification.* 13D45, 13C14.

Key words and phrases. Top local cohomology, Depth, Tensor products, Matlis duality, Cohen-Macaulayness.

The author is supported in part by a grant from IPM (No. 91130026).

$D(H^d_{\mathfrak{a}}(R))$. In particular, in the case R is complete and $\operatorname{pd}_{R/Q}(K_{R/Q}) < \infty$ for a certain ideal Q of R we give the necessary and sufficient conditions for Cohen-Macaulayness of $D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))$ (see Theorem 4.2). We denote by $\operatorname{pd}_R(M)$ resp. K_M the projective dimension resp. the canonical module of a finite R-module M (if it exists).

2. PRELIMINARIES

Let $(R, \mathfrak{m})$ be a local ring of dimension d. We denote by $E_R(R/\mathfrak{m})$ the injective hull of the residue field $R/\mathfrak{m} = k$. The Matlis duality functor $\operatorname{Hom}_R(-, E_R(R/\mathfrak{m}))$ is denoted by $D(-)$. We state the following lemmas which we need in our proofs.

Lemma 2.1. *Let $(R, \mathfrak{m})$ be a complete local ring. Suppose that M and N are either Artinian or finitely generated R-modules. Then we have the following isomorphism*

$$\operatorname{Hom}(M, N) \cong \operatorname{Hom}(D(N), D(M)).$$

Proof. Since $N \cong D(D(N))$ so by [R, Theorem 2.11] we get the claim as follows:

$$\begin{aligned} \operatorname{Hom}_R(M, N) &\cong \operatorname{Hom}_R(M, D(D(N))) \\ &\cong \operatorname{Hom}_R(D(N), D(M)). \end{aligned}$$

□

Lemma 2.2. *Let $(R, \mathfrak{m})$ be a complete local ring. Let M and N be Artinian R-modules. Then we have the following isomorphism*

$$D(N) \otimes_R M \cong D(\operatorname{Hom}_R(M, N)).$$

Proof. It should be noted that, by Matlis duality $D(N)$ is a finitely generated R-modules. By Hom $-\otimes$-adjointness we can prove the claim as follows:

$$\begin{aligned} D(N) \otimes_R M &\cong D(N) \otimes_R D(D(M)) \\ &\cong D(N) \otimes_R \operatorname{Hom}_R(D(M), E(R/\mathfrak{m})) \\ &\cong \operatorname{Hom}_R(\operatorname{Hom}_R(D(N), D(M)), E(R/\mathfrak{m})) \qquad \text{(cf. [Br-Sh, 10.2.16])} \\ &\cong \operatorname{Hom}_R(\operatorname{Hom}_R(M, N), E(R/\mathfrak{m})). \end{aligned}$$

The last isomorphism follows by 2.1. □

We need the definition of the canonical module of a finitely generated R-module M. To this end let $(R, \mathfrak{m})$ be the epimorphic image of a local Gorenstein ring $(S, \mathfrak{n})$ and $n = \dim S$.

Definition 2.3. (A) For a finitely generated R-module M we consider

$$K(M) = \operatorname{Hom}_R(H^t_{\mathfrak{m}}(M), E_R(R/\mathfrak{m})),\ t = \dim M.$$

Because $H^t_{\mathfrak{m}}(M)$ is an Artinian R-module it turns out that $K(M)$ is a finitely generated $\hat{R}$-module.
(B) This is closely related to the notion of the canonical module K_M of a finitely generated R-module M. To this end we have to assume that $(R, \mathfrak{m})$ is the epimorphic image of a local Gorenstein ring $(S, \mathfrak{n})$ with $n = \dim S$. Then $K_M = \operatorname{Ext}^{n-t}_S(M, S)$ is called the canonical module of M.

By virtue of the Cohen's Structure Theorem and (A) there is an isomorphism $K(M) \simeq K_{\hat{M}}$. By the Local Duality Theorem (see [G]) there are the following isomorphisms

$$\operatorname{Hom}_R(H^t_{\mathfrak{m}}(M), E_R(R/\mathfrak{m})) \simeq K(M) \simeq K_{\hat{M}}.$$

Here $\hat{R}$ resp. $\hat{M}$ denote the $\mathfrak{m}$-adic completion of R resp. of M.

In the following, we summarize a few results about the last non-vanishing local cohomology module $H^d_{\mathfrak{a}}(R)$. To this end note that a finite module M over a Noetherian ring R satisfies Serre's condition S_n if $\operatorname{depth} M_{\mathfrak{p}} \geq \min\{n, \dim M_{\mathfrak{p}}\}$ for all $\mathfrak{p} \in \operatorname{Supp} M$.

Lemma 2.4. *Let $(R, \mathfrak{m})$ be a complete local ring of dimension d and $H^d_{\mathfrak{a}}(R) \neq 0$.*

(a) *$H^d_{\mathfrak{a}}(R)$ is an Artinian R-module.*
(b) *$H^d_{\mathfrak{a}}(R) \cong H^d_{\mathfrak{m}}(R/Q_{\mathfrak{a}}(R))$, where*

$$Q_{\mathfrak{a}}(R) = \bigcap\{\mathfrak{q} : \mathfrak{q} \textit{ primary component of } R,\ \dim R/\mathfrak{q} = d \textit{ and } \operatorname{Rad}(\mathfrak{a} + \mathfrak{q}) = \mathfrak{m}\}.$$

(c) *$D(H^d_{\mathfrak{a}}(R)) \cong K_{R/Q_{\mathfrak{a}}(R)}$ is a finitely generated R-module.*
(d) *$\operatorname{Ann}_R H^d_{\mathfrak{a}}(R) = Q_{\mathfrak{a}}(R)$.*
(e) *$\dim D(H^d_{\mathfrak{a}}(R)) = d$.*
(f) *The module $D(H^d_{\mathfrak{a}}(R))$ satisfies Serre's condition S_2.*
(g) *If $R/Q_{\mathfrak{a}}(R)$ is a Cohen-Macaulay ring, then the module $D(H^d_{\mathfrak{a}}(R))$ is Cohen-Macaulay.*

Proof. For the proof of (a), see [Br-Sh, Theorem 7.1.6]. For the proof of (b), (c), (d) and (g) we refer to [E-Sch, 3.3, 4.1(c), 4.2(a) and 4.3]. Part (e) follows by (b) and (d). The prove of (f) is a consequence of (c) and [Sch1, Lemma 1.9]. ☐

As a technical tool, in Section three, we use the so-called formal local cohomology modules. We denote by $\mathfrak{F}^i_{\mathfrak{a}}(M) := \varprojlim_n H^i_{\mathfrak{m}}(M/\mathfrak{a}^n M)$ the ith formal local cohomology of M with respect to $\mathfrak{a}$. see [Sch2] for more information and also [E].

Lemma 2.5. *(see* [Sch2, Remark 3.6 and Theorem 4.5]*) Let $(R, \mathfrak{m})$ be a local ring.*

(a) *If M is a finitely generated R-module, then*

$$\dim_R M/\mathfrak{a}M = \sup\{i \in \mathbb{Z} : \mathfrak{F}^i_{\mathfrak{a}}(M) \neq 0\}.$$

(b) *If the local ring $(R, \mathfrak{m})$ is a quotient of a local Gorenstein ring $(S, \mathfrak{n})$, Then*

$$\mathfrak{F}^i_{\mathfrak{a}}(M) \cong \operatorname{Hom}_R(H^{n-i}_{\mathfrak{a}S}(M, S), E), i \in \mathbb{Z},$$

where M is considered as an S-module, $n = \dim S$ and E denotes the injective hull of the residue field $R/\mathfrak{m}$.

It should be noted that $H^i_{\mathfrak{a}}(M, N) = \varinjlim_n \operatorname{Ext}^i_R(M/\mathfrak{a}^n M, N)$ is known as the ith generalized local cohomology module provided that M and N are finitely generated R-modules. In case $R = M$, the module $H^i_{\mathfrak{a}}(M, N)$ is the same as the ordinary local cohomology module $H^i_{\mathfrak{a}}(N)$.

3. ON THE DEPTH

Let $(R, \mathfrak{m})$ be a complete local ring of dimension d and $\mathfrak{a}$ be an ideal of R. It is a well-known result that $D(H^d_{\mathfrak{a}}(R))$ is a Cohen-Macaulay module of dimension d for $d \leq 2$ (see for example 2.4(f)). In this section we examine the depth of Matlis dual of top local cohomology modules.

Theorem 3.1. *Let $(R, \mathfrak{m})$ be a complete local ring of dimension d. Let $H^d_{\mathfrak{a}}(R)$ be nonzero. Then*

$$\operatorname{depth}(D(H^d_{\mathfrak{a}}(R)) \geq \min\{d, 2\}.$$

Proof. We prove the claim by induction on d. For $d \leq 2$, the claim is clear. Assume that $d \geq 3$.

First note that by virtue of Lemma 2.4 we have $H^d_{\mathfrak{a}}(R) \cong H^d_{\mathfrak{m}}(R/Q)$, where $Q := Q_{\mathfrak{a}}(R)$. By the definition of Q we may deduce that $\mathfrak{m} \notin \operatorname{Ass} R/Q$ and so $\operatorname{depth} R/Q > 0$. Hence, there exists an element $x \in \mathfrak{m}$ which is nonzero divisor on R/Q. From the short exact sequence

$$0 \to R/Q \xrightarrow{x} R/Q \to R/(x, Q) \to 0$$

we get the following long exact sequence

$$0 \to D(H^d_{\mathfrak{m}}(R/Q)) \xrightarrow{x} D(H^d_{\mathfrak{m}}(R/Q)) \to D(H^{d-1}_{\mathfrak{m}}(R/(x, Q)) \to \cdots .$$

Since $\dim(R/(x, Q)) = d - 1$, so by induction hypothesis one can see that

$$\operatorname{depth} D(H^{d-1}_{\mathfrak{m}}(R/(x, Q)) \geq 2.$$

Hence, there are $u, v \in \mathfrak{m} \setminus \bigcup_{\mathfrak{p} \in \operatorname{Ass}(D(H^d_{\mathfrak{m}}(R/Q))/xD(H^d_{\mathfrak{m}}(R/Q)))} \mathfrak{p}$. Thus it follows that

$$\operatorname{depth}(D(H^d_{\mathfrak{m}}(R/Q)) \geq 2.$$

□

According to [H, Corollary 1.1.4] the $\operatorname{grade}(\mathfrak{a}, D(H^d_{\mathfrak{a}}(R)))$ plays a significant role to see whether $\mathfrak{a}$ is a set-theoretic complete intersection. In this direction, we try to find an explicit relation between $\operatorname{depth} D(H^d_{\mathfrak{a}}(R))$ and $\operatorname{grade}(\mathfrak{a}, D(H^d_{\mathfrak{a}}(R)))$ (see Theorem 3.2). To this end we denote by $\operatorname{grade}_{\mathfrak{a}}(M, N)$ the least integer i such that $H^i_{\mathfrak{a}}(M, N)$ is nonzero for finitely generated R-modules M and N.

Theorem 3.2. *Let $\mathfrak{a}$ be an ideal of a local ring $(R, \mathfrak{m})$. Let M and N be finitely generated R-modules. Then*

$$\operatorname{grade}_{\mathfrak{a}}(M, N) \geq \max\{0, \operatorname{depth} N - \dim M/\mathfrak{a}M\}.$$

In order to prove Theorem 3.2 we need some preparations.

Remark 3.3. Let $(R, \mathfrak{m})$ be a local ring which is a factor ring of a Gorenstein local ring $(S, \mathfrak{n})$. In the light of Lemma 2.5 we have

$$\begin{aligned} \dim M/\mathfrak{a}M &= \sup\{i : \mathfrak{F}^i_{\mathfrak{a}}(M) \neq 0\} \\ &= \sup\{i : H^{\dim S - i}_{\mathfrak{a}S}(M, S) \neq 0\} \\ &= \sup\{\dim S - j : H^j_{\mathfrak{a}S}(M, S) \neq 0\} \\ &= \dim S - \inf\{i : H^i_{\mathfrak{a}S}(M, S) \neq 0\} \\ &= \dim S - \operatorname{grade}_{\mathfrak{a}}(M, S), \end{aligned}$$

where

$$\dim M/\mathfrak{a}M = \dim S - \operatorname{grade}(\mathfrak{a} + \operatorname{Ann}_S M).$$

See [B, Proposition 5.5].

Lemma 3.4. *Let M and N be two finitely generated R-modules and $\mathfrak{a}$ be an ideal of R. Then*

$$\operatorname{grade}_{\mathfrak{a}}(M, N) \geq \operatorname{grade}_{\mathfrak{a}}(M, R) - \operatorname{pd}_R N,$$

provided that $\operatorname{pd}_R N < \infty$ *.*

Proof. We argue by induction on $n := \operatorname{pd}_R N$.

Put $n = 0$ so N is a finite free R-module of the form $N \cong \oplus_t R$, where t is an integer. Hence, the isomorphisms

$$H^i_{\mathfrak{a}}(M, N) \cong H^i_{\mathfrak{a}}(M, \oplus_t R) \cong \oplus_t H^i_{\mathfrak{a}}(M, R)$$

hold. Clearly $\operatorname{grade}_{\mathfrak{a}}(M, N) \geq \operatorname{grade}_{\mathfrak{a}}(M, R)$.

Next, suppose that $n > 0$ and the claim holds for $n - 1$. There is a finite free R-module F such that

$$0 \to L \to F \to N \to 0$$

which induces $\operatorname{pd}_R L = n - 1$. By induction hypothesis $\operatorname{grade}_{\mathfrak{a}}(M, L) \geq \operatorname{grade}_{\mathfrak{a}}(M, R) - n + 1$. If $i < \operatorname{grade}_{\mathfrak{a}}(M, R) - n$, then it follows from the long exact sequence

$$\cdots \to H^i_{\mathfrak{a}}(M, F) \to H^i_{\mathfrak{a}}(M, N) \to H^{i+1}_{\mathfrak{a}}(M, L) \to \cdots$$

that $H^i_{\mathfrak{a}}(M, N)$ is zero. Hence, $\operatorname{grade}_{\mathfrak{a}}(M, N) \geq \operatorname{grade}_{\mathfrak{a}}(M, R) - \operatorname{pd}_R N$. □

Now is the time to give the proof of Theorem 3.2 as follows:

Proof of Theorem 3.2. First note that by passing to the completion we may assume that R is a complete local ring. Then, by Cohen's structure Theorem R is a factor ring of a regular local ring $(T, \mathfrak{n})$. In the light of Independence Theorem we may assume that $\mathfrak{F}^i_{\mathfrak{a}}(M) \cong \mathfrak{F}^i_{\mathfrak{b}}(M)$, where $\mathfrak{b} := \mathfrak{a} \cap T$.

Using Remark 3.3, we have

$$\dim M/\mathfrak{a}M = \dim T - \operatorname{grade}_{\mathfrak{a}}(M, T).$$

So by Lemma 3.4 and Auslander-Buchsbaum Theorem we prove the claim as follows:

$$\begin{aligned} \dim M/\mathfrak{a}M &= \dim T - \operatorname{grade}_{\mathfrak{a}}(M, T) \\ &\geq \dim T - \operatorname{grade}_{\mathfrak{a}}(M, N) - \operatorname{pd}_T N \\ &= \operatorname{depth} N - \operatorname{grade}_{\mathfrak{a}}(M, N). \end{aligned}$$

□

Assume that $\mathfrak{b}$ is an ideal of R of dimension at most 2 and $d \geq 3$. By virtue of Theorem 3.2 and 3.1 we have

$$\operatorname{grade}(\mathfrak{b}, D(H^d_{\mathfrak{a}}(R))) + \dim R/\mathfrak{b} \geq \operatorname{depth}(D(H^d_{\mathfrak{a}}(R))) \geq 2,$$

which is an upper bound for $\operatorname{depth}(D(H^d_{\mathfrak{a}}(R)))$.

Remark 3.5. Let $(R, \mathfrak{m})$ be a local ring of dimension d.

(a) Let x be a parameter element of R. Consider $0 :_R < x > = \bigcup_{n\geq 1} 0 :_R x^n$ and following exact sequence

$$0 \to 0 :_R < x > \to R \to R/0 :_R < x > \to 0.$$

One can see that $\dim(0 :_R < x >) < d$ and $\dim R/(0 :_R < x >) = d$. Now apply $H^d_{\mathfrak{a}}(-)$ to the above short exact sequence to get the following exact sequence

$$H^d_{\mathfrak{a}}(0 :_R < x >) \to H^d_{\mathfrak{a}}(R) \to H^d_{\mathfrak{a}}(R/0 :_R < x >) \to 0.$$

Hence, by Grothendieck's vanishing Theorem, we have the isomorphism $H^d_{\mathfrak{a}}(R) \cong H^d_{\mathfrak{a}}(R/0 :_R < x >)$.

(b) Now we give conditions that Cohen-Macaulayness of $D(H^d_{\mathfrak{a}}(R))$ fails. Let $(R, \mathfrak{m})$ be a complete local ring, $d \geq 3$ and $\mathfrak{b}$ be a one-dimensional ideal containing a regular element x of R (part one guarantees the existence of a regular element of R). Then, it implies the following short exact sequence

$$0 \to R \xrightarrow{x} R \to R/xR \to 0.$$

Thus we have the following long exact sequence

$$\cdots \to H^{d-1}_{\mathfrak{a}}(R/xR) \to H^d_{\mathfrak{a}}(R) \xrightarrow{x} H^d_{\mathfrak{a}}(R) \to 0.$$

By applying $D(-)$ we get

$$0 \to D(H^d_{\mathfrak{a}}(R)) \xrightarrow{x} D(H^d_{\mathfrak{a}}(R)) \to D(H^{d-1}_{\mathfrak{a}}(R/xR)) \to \cdots.$$

Now it yields that x is a $D(H^d_{\mathfrak{a}}(R))$-regular element. So $\operatorname{grade}(\mathfrak{b}, D(H^d_{\mathfrak{a}}(R))) > 0$. Suppose that $\mathfrak{b}$ does not contain a $D(H^d_{\mathfrak{a}}(R))/xD(H^d_{\mathfrak{a}}(R))$-regular element. Hence, [Bru-Her, 1.2.3 and 1.2.4] imply that $\operatorname{Ext}^1_R(R/\mathfrak{b}, D(H^d_{\mathfrak{a}}(R))) \neq 0$ and then, $\operatorname{grade}(\mathfrak{b}, D(H^d_{\mathfrak{a}}(R))) = 1$. Therefore, in the view of Theorem 3.1 and Theorem 3.2 we have

$$2 \leq \operatorname{depth}(D(H^d_{\mathfrak{a}}(R)) \leq \operatorname{grade}(\mathfrak{b}, D(H^d_{\mathfrak{a}}(R))) + \dim R/\mathfrak{b} = 2.$$

It shows that with the above assumptions $D(H^d_{\mathfrak{a}}(R))$ is not Cohen-Macaulay for $d \geq 3$.

4. TENSOR PRODUCTS

Throughout this section, $(R, \mathfrak{m})$ is a local ring of dimension d. We assume that $H^d_{\mathfrak{a}}(R) \neq 0$ and $Q := Q_{\mathfrak{a}}(R)$. In recent research there is an interest in endomorphism rings of certain local cohomology modules $H^i_{\mathfrak{a}}(R)$. This was done in the case of $i = d$ by the author and Schenzel (see [E-Sch]) and some conditions for Cohen-Macaulayness of $\operatorname{Hom}_R(H^d_{\mathfrak{a}}(R), H^d_{\mathfrak{a}}(R))$ have been investigated. For the sake of completeness, we state the following result.

Remark 4.1. Let $\mathfrak{a}$ be an ideal of a complete local ring $(R, \mathfrak{m})$. Then,

$$\operatorname{depth}_R(\operatorname{Hom}_R(H^d_{\mathfrak{a}}(R), H^d_{\mathfrak{a}}(R))) \geq \min\{2, d\}.$$

To this end note that $\operatorname{Hom}_R(H^d_{\mathfrak{a}}(R), H^d_{\mathfrak{a}}(R)) \cong \operatorname{Hom}_R(D(H^d_{\mathfrak{a}}(R)), D(H^d_{\mathfrak{a}}(R)))$ (see 2.1). Hence, the claim is clear by virtue of [Bru-Her, Exer. 1.4.19] and Theorem 3.1.

The next part of our work is to investigate the tensor product of top local cohomology modules and their Matlis duals.

Theorem 4.2. *Let $(R, \mathfrak{m})$ be a complete local ring of dimension d. Then, the following statements hold:*

(a) $H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R)$ *is of finite length.*

(b) *Assume that R/Q is a Gorenstein ring. Then,*

$$H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R) = 0.$$

(c) *The ideal Q is contained in $\operatorname{Ann}_R(D(H^d_{\mathfrak{a}}(R)) \otimes_R D(H^d_{\mathfrak{a}}(R))$ and the Krull dimension of $D(H^d_{\mathfrak{a}}(R)) \otimes_R D(H^d_{\mathfrak{a}}(R))$ is equal to d.*

(d) *Suppose that $\operatorname{pd}_{R/Q}(K_{R/Q}) < \infty$ then, the following statements are equivalent :*

  (1) *R/Q is a Cohen-Macaulay ring.*

  (2)

$$\operatorname{Tor}^{R/Q}_i(D(H^d_{\mathfrak{a}}(R)), D(H^d_{\mathfrak{a}}(R))) = 0 \ \textit{ for all } \ i \geq 1$$

  and

$$D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R)) \quad \textit{is a Cohen-Macaulay R/Q-module.}$$

(e) *Assume that R/Q satisfies S_2 situation. Then, there is an isomorphism*

$$H^d_{\mathfrak{a}}(R) \otimes_R D(H^d_{\mathfrak{a}}(R)) \cong E_{R/Q}(R/\mathfrak{m}).$$

Proof. (a) As $H^d_{\mathfrak{a}}(R)$ is an Artinian R-module, then it follows by [F-H, Proposition 6.1] that $H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R)$ is of finite length.

(b) Since $H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R)$ is of finite length so by Hom $-\otimes$-adjointness we have

$$\begin{aligned} H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R) &\cong D(D(H^d_{\mathfrak{a}}(R) \otimes_R H^d_{\mathfrak{a}}(R))) \\ &\cong D(\operatorname{Hom}_R(H^d_{\mathfrak{a}}(R), D(H^d_{\mathfrak{a}}(R))) \\ &\cong D(\operatorname{Hom}_R(H^d_{\mathfrak{a}}(R), K_{R/Q})). \end{aligned}$$

As R/Q is a Gorenstein ring of positive dimension then, the later module is zero. To this end note that the last isomorphism follows by Lemma 2.4.

(c) It is clear that

$$\operatorname{Ann}_R(D(H^d_{\mathfrak{a}}(R)) \otimes_R D(H^d_{\mathfrak{a}}(R)) \supseteq \operatorname{Ann}_R(D(H^d_{\mathfrak{a}}(R))) = Q$$

(see 2.4) and

$$\begin{aligned} \dim_R(D(H^d_{\mathfrak{a}}(R)) \otimes_R D(H^d_{\mathfrak{a}}(R))) &= \dim_R(\operatorname{Supp}(D(H^d_{\mathfrak{a}}(R)) \otimes_R D(H^d_{\mathfrak{a}}(R)))) \\ &= \dim_R D(H^d_{\mathfrak{a}}(R)) = d \end{aligned}$$

(see 2.4).

(d) It should be noted that for a finite module M which has R-module and R/Q-module structures then, it is clear that $\operatorname{depth}_R M = \operatorname{depth}_{R/Q} M$. Moreover, by virtue of 2.4 it follows that $D(H^d_{\mathfrak{a}}(R)) \cong K_{R/Q}$.

In order to prove that 1 implies 2, take into account that if R/Q is a Cohen-Macaulay ring, then $D(H^d_{\mathfrak{a}}(R))$ is a Cohen-Macaulay module (see 2.4). Thus by Auslander-Buchsbaum Theorem we deduce that $K_{R/Q}$ is a free R/Q-module. Hence,

$$\operatorname{Tor}^{R/Q}_i(D(H^d_{\mathfrak{a}}(R)), D(H^d_{\mathfrak{a}}(R))) = 0, \text{ for all } i \geq 1.$$

Now using [Hu-W, Lemma 2.2] we have

$$\operatorname{depth}(D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))) = d.$$

Thus $D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))$ is a Cohen-Macaulay R/Q-module.

To show that (2) implies (1), we first use the hypothesis and [Hu-W, Lemma 2.2]. Then we have

$$2 \operatorname{depth} D(H^d_{\mathfrak{a}}(R)) = \operatorname{depth} R/Q + \operatorname{depth}(D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))). \tag{4.1}$$

As $\operatorname{pd}_{R/Q}(K_{R/Q}) < \infty$, then by Auslander-Buchsbaum Theorem it follows that

$$\operatorname{depth} D(H^d_{\mathfrak{a}}(R)) \leq \operatorname{depth} R/Q.$$

Since $D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))$ is a Cohen-Macaulay R/Q-module, then in the light of the equality (4.1) we prove the claim as follows:

$$\begin{aligned} \operatorname{depth} R/Q &\geq \operatorname{depth}(D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))) \\ &= \dim_{R/Q}(D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))) \\ &= \dim R/Q. \end{aligned}$$

(e) As $H^d_{\mathfrak{a}}(R)$ is an Artinian R-module so $D(H^d_{\mathfrak{a}}(R))$ is finite. Hence,

$$\begin{aligned} H^d_{\mathfrak{a}}(R) \otimes_R D(H^d_{\mathfrak{a}}(R)) &\cong D(\operatorname{Hom}_R(H^d_{\mathfrak{a}}(R), H^d_{\mathfrak{a}}(R))) \\ &\cong D(R/Q) \\ &\cong E_{R/Q}(R/\mathfrak{m}). \end{aligned}$$

To this end note that the first isomorphism follows by Lemma 2.2 and the second one follows by [E-Sch, Theorem 4.2].

□

Let M and N be nonzero finitely generated modules over a local ring $(R, \mathfrak{m})$. M and N satisfy the depth formula provided that

$$\operatorname{depth} M + \operatorname{depth} N = \operatorname{depth} R + \operatorname{depth}(M \otimes_R N).$$

A necessary condition for the depth formula to hold is that $\operatorname{depth} M + \operatorname{depth} N \geq \operatorname{depth} R$ (see [Hu-W, pp. 163]). As an application of 3.1 we deduce the following result:

Corollary 4.3. *Let $(R, \mathfrak{m})$ be a d-dimensional complete local ring. In case $d = 3$ or 4, then we have*

$$\operatorname{depth}(D(H^d_{\mathfrak{a}}(R) \otimes_R D(H^d_{\mathfrak{a}}(R)) \geq 4 - \operatorname{depth} R.$$

Corollary 4.4. *Let $(R, \mathfrak{m})$ be a complete local ring with $d \geq 4$. Suppose that $\operatorname{pd}_{R/Q}(K_{R/Q}) < \infty$ and $\operatorname{Tor}_i^{R/Q}(D(H^d_{\mathfrak{a}}(R)), D(H^d_{\mathfrak{a}}(R))) = 0$ for all $i > 0$. Then,*

(a) $\operatorname{depth}(D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))) \geq 4 - \operatorname{depth} R/Q$,

(b) $\operatorname{pd}_{R/Q}(D(H^d_{\mathfrak{a}}(R)) \otimes_{R/Q} D(H^d_{\mathfrak{a}}(R))) \leq 2 \operatorname{depth} R/Q - 4$.

Proof. By Lemma 2.4 it follows that $D(H^d_{\mathfrak{a}}(R)) \cong K_{R/Q}$ is a finitely generated module. Part (a) is a consequence of 3.1 and [A, Theorem 1.2]. Part (b) follows by (a) and the Auslander-Buchsbaum Theorem. □

REFERENCES

[A] M. Auslander, *Modules over unramified regular local rings*, Ill. J. Math. **5** (1961), 631–647.

[B] M. H. Bijan-Zadeh, *A common generalization of local cohomology theories*, Glasgow Math. J. **21**, 173–181, (1980).

[Br-Sh] M. Brodmann and R.Y. Sharp, *Local cohomology: an algebraic introduction with geometric applications*, Cambridge Univ. Press, **60**, Cambridge, (1998).

[Bru-Her] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge Univ. Press, (1993).

[E] M. Eghbali, *On Artinianness of formal local cohomology, colocalization and coassociated primes*. Math. Scand. , to appear. arXiv:1011.4141 [math.AC].

[E-Sch] M. Eghbali and P. Schenzel, *On an endomorphism ring of local cohomology* , Comm. Alg. **40**: 4295–4305, (2012).

[F-H] C. Faith and D. Herbera, *Endomorphism rings and tensor products of linearly compact modules*, Comm. Alg. **25**(4), (1997), 1215–1255.

[G] A. Grothendieck, *Local cohomology*, Notes by R. Hartshorne, Lect. Notes in Math. **20**, Springer, 1966.

[H] M. Hellus, *Local Cohomology and Matlis duality*, Habilitationsschrift, Leipzig (2006).

[H-Sch] M. Hellus and P. Schenzel, *On cohomologically complete intersections*, J. Algebra **320** (2008) 3733–3748.

[H-St] M. Hellus and J. Stückrad, *On endomorphism rings of local cohomology modules*, Proc. A.M.S Vol. **136**(7), (2008), (2333–2341).

[Hu-W] C. Huneke and R. Wiegand, *Tensor products of modules, rigidity and local cohomology* , Math. Scand. **81** (1997), 161–183.

[R] J. Rotman, *An Introduction to Homological Algebra*, Academic Press, Orlando, FL, 1979.

[Sch1] P. Schenzel, *On the use of local cohomology in algebra and geometry*, In: J. Elias, J.M. Giral, R.M. Miró-Roig, S. Zarzuela (Eds.), Six Lectures in Commutative Algebra, Proceed. Summer School on Commutative Algebra at Centre de Recerca Matemtica, Progr. Math., Vol. **166**, Birkhäuser, 1998, 241-292.

[Sch2] P. Schenzel, *On formal local cohomology and connectedness*, J. Algebra, **315**(2), (2007), 894–923.

[Sch3] P. Schenzel, *On endomorphism rings and dimensions of local cohomology modules* , Proc. A.M.S Vol. **137**(4), (2009), (1315–1322).

M. EGHBALI , SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P. O. BOX: 19395-5746, TEHRAN-IRAN.

E-mail address: `m.eghbali@yahoo.com`